

# IMMUNIZATION SCHEDULE

## Recommended immunizations & tests at each check up

Physical exams are recommended at 2 weeks, 2, 4, 6,9,12, 15,18, 24 & 30 months and annually from 3 years on. Physical exams are necessary to provide medical information required by some camp and school forms. **No more than 3 injections with one oral will be given at any one visit.**

State law requires the following minimum doses: **5** Dtap, **4** IPV, **4** HIB, **3** Hep B, **2** MMR, **1** Varicella

2 wks/1 month	No scheduled tests at this check up					
2 months	DTap (Pentacel)	Hib (Pediarix)	Pevnar	IPV	Hep B	Rotavirus
4 months	DTap (Pentacel)	Hib (Pediarix)	Pevnar	IPV	Hep B	Rotavirus
6 months	DTap (Pentacel)	Hib (Pediarix)	Pevnar	IPV	Hep B	Rotavirus
9 months	No scheduled tests at this check up					
12 months	Pevnar	Varicella*				
15 months	MMR	Hib	Hgb	Lead Level (all Medicaid or if at risk)		
18 months	DTap					
2 years	No scheduled tests at this check up					
3 years	Vision (must attempt)		Lead Level (all Medicaid or if at risk)			
4 years	DTap	MMR	Vision/Hearing			
5 years	IPV	Varicella*	Vision/Hearing			
9-18 years	Hgb (once, after start of menses for females)					
11-13 years	Menactra	Tdap	HPV (Gardasil)			
16-21 years	Booster Menactra (5 yrs after first dose)					

\*Two doses of the varicella (chicken pox) vaccine are recommended by the American Academy of Pediatrics Committee on Infectious Diseases: 1<sup>st</sup> dose at 12 to 15 months of age 2<sup>nd</sup> dose between 4 and 6 years of age

### Who should get Tdap?

- Ages 7-10 who has not had any Pertussis vaccine or has not finished the series
- Ages 65 and older, especially if in contact with newborn/infants

Tdap can be given regardless of the time since previous tetanus booster.

### HPV (Gardasil) Spacing

- Dose 2: 2 months after dose 1
- Dose 3: 6 months after dose 1

\*\*Do not give HPV with Tdap or Menactra

DTap (IM)	Rotarix (PO)
Hib (IM)	MMR (SQ)
Menactra (IM)	Tdap (IM)
IPV (SQ)	Varicella (SQ)
Hep A (IM)	Pevnar (IM)
Hep B (IM)	HPV (IM)

Pentacel (IM) = DTap/IPV/Hib  
Pediarix (IM) = DTap/IPV/Hep B

Do not give flu vaccine with DTap, Tdap, HPV, MMR, or Varicella

### When to give Menactra (MCV4) boosters

- If received initial dose at age 11-12, booster at 16 or up to 5 years after initial dose
- If received initial dose at age 13-15, booster at age 16-18