

Benadryl can help reduce your child's symptoms during an allergic reaction. Benadryl, which is also available as generic **Diphenhydramine**, is an over-the-counter antihistamine for use during allergic reactions. Benadryl dosing is based on weight, **not age**. The following chart can help determine the correct dose for your child **based on your child's weight**, but is no substitute for your doctor's advice. Therefore, **you should always call the pediatrician if you have any questions.** Benadryl is available in liquid, chewable, and tablet forms. Chewable form may be a choking hazard and are not recommended for children under 18 months.

## Use the following tips to keep your child safe and avoid over dosage:

- Benadryl is also a sedative and care must be taken not to overdose your child.
- Give every 6 hours as needed.
- Do not exceed 4 doses in 24 hours.
- Do not give Benadryl to children under 1 year without consulting with your doctor first.
- Always read label first, since different medicines have difference concentrations
- Always use measuring device (dropper or measuring cup) that comes with the medication or you could risk overdosing. Never use droppers from different medications or spoons from the kitchen.

## Remember to call the pediatrician for:

- Children under 1 year old.
- If your child is flushed, coughing, drooling or having difficulty breathing call 911 and proceed to the nearest emergency room.

Following are the dosage recommendations:

Child's Weight	Liquid Benadryl	Benadryl Chewable	Benadryl Tablet
	(12.5 mg/ 5ml)	Tablet	(25 mg)
		(12.5 mg)	
17-21 lbs	.75 tsp (3.75 ml)	X	Х
22-32 lbs	1 tsp (5 ml)	1 chewable	Х
33-42 lbs	1.5 tsp (7.5 ml)	1.5 chewable	Х
43-53 lbs	2 tsp (10 ml)	Х	1 tablet
54-64 lbs	2.5 tsp (12.5 ml)	2 chewables	1 tablet
65-75 lbs	3 tsp (15 ml)	3 chewables	1 tablet
76-86 lbs	3.5 tsp (17.5 ml)	4 chewables	1 tablet
>86 lbs	4 tsp (20 ml)	4 chewables	2 tablets